

(19) World Intellectual Property  
Organization  
International Bureau



(43) International Publication Date  
1 September 2005 (01.09.2005)

PCT

(10) International Publication Number  
**WO 2005/079758 A1**

(51) International Patent Classification<sup>7</sup>: **A61K 9/113**,  
31/05

TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM,  
ZW.

(21) International Application Number:  
PCT/US2005/004533

(84) Designated States (unless otherwise indicated, for every  
kind of regional protection available): ARIPO (BW, GH,  
GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM,  
ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM),  
European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI,  
FR, GB, GR, HU, IE, IS, IT, LT, LU, MC, NL, PL, PT, RO,  
SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN,  
GQ, GW, ML, MR, NE, SN, TD, TG).

(22) International Filing Date: 14 February 2005 (14.02.2005)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:  
60/544,604 13 February 2004 (13.02.2004) US

**Declarations under Rule 4.17:**

— as to applicant's entitlement to apply for and be granted  
a patent (Rule 4.17(ii)) for the following designations AE,  
AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ,  
CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE,  
EG, ES, FI, GB, GD, GE, GH, GM, GR, HU, ID, IL, IN, IS,  
JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA,  
MD, MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM,  
PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SM, SY,  
TJ, TM, TN, TR, TT, TZ, UA, UG, UZ, VC, VN, YU, ZA,  
ZM, ZW, ARIPO patent (BW, GH, GM, KE, LS, MW, MZ,  
NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM,  
AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT,  
BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR,  
HU, IE, IS, IT, LT, LU, MC, NL, PL, PT, RO, SE, SI, SK,  
TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ,  
GW, ML, MR, NE, SN, TD, TG)

— of inventorship (Rule 4.17(iv)) for US only

(71) Applicant (for all designated States except US):  
**BIOAVAILABILITY, INC.** [US/US]; 10743 Civic  
Center Drive, Rancho Cucamonga, CA 91730 (US).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **HARRIS, Steven, B.**  
[US/US]; 10743 Civic Center Drive, Rancho Cucamonga,  
CA 91730 (US). **HUANG, Nick, J.** [CN/US]; 10743 Civic  
Center Drive, Rancho Cucamonga, CA 91730 (US).

(74) Agent: **HENDRICKSON, Jay, P.**; 1010 B Street, Suite  
319, San Rafael, CA 94901 (US).

(81) Designated States (unless otherwise indicated, for every  
kind of national protection available): AE, AG, AL, AM,  
AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN,  
CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI,  
GB, GD, GE, GH, GM, GR, HU, ID, IL, IN, IS, JP, KE,  
KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD,  
MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG,  
PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM,

**Published:**

— with international search report

For two-letter codes and other abbreviations, refer to the "Guid-  
ance Notes on Codes and Abbreviations" appearing at the begin-  
ning of each regular issue of the PCT Gazette.

(54) Title: A MICROEMULSION PREPARATION OF HIGH CONCENTRATION PROPOFOL FOR ANESTHETIC USES

(57) Abstract: The invention provides a method and a composition for enhancing the dissolution and bioavailable properties of propofol (2, 6 diisopropyl phenol) for use as an intravenously administered anesthetic in mammals. The method produces a self-microemulsifiable emulsion base composition that is utilized in the production of a water-based microemulsion preparation. In a preferred two (2) component base composition, the base composition consists of: a surfactant, containing polyethylene glycol; and liquid propofol. The microemulsion is prepared by mixing the base composition with a carrier liquid, which results in the formation of a microemulsion containing concentrations of propofol of up to about 4% by weight of propofol to the volume of the microemulsion. In a four (4) component base composition, the base composition consists of: a surfactant, containing polyethylene glycol; liquid propofol; a water-immiscible solvent; and ethanol. The microemulsion is prepared by mixing the base composition with a carrier liquid, which results in the formation of a microemulsion containing concentrations of propofol of up to about 10% by weight of propofol to the volume of the microemulsion.



WO 2005/079758 A1